Hydrazine Kit

K-5011/R-5011 ULR CHEMets® Kit

0 - 50 ppb

K-5005/R-5005 CHEMets® Kit:


0 - 0.5 ppm


Safety Information


Read ŠDS (available at www.chemetrics.com) before performing this test procedure. Wear safety glasses and protective gloves.

Test Procedure

- 1. Fill the sample cup to the 25 mL mark with the sample to be tested (fig. 1).
- 2. Place the CHEMet ampoule, tip first, into the sample cup. Snap the tip. The ampoule will fill leaving a bubble for mixing (fig. 2).
- To mix the ampoule, invert it several times, allowing the bubble to travel from end to end.
- 4. Dry the ampoule and wait **10 minutes** for color development.
- Obtain a test result by placing the ampoule, flat end first, into the comparator. Hold the comparator up toward a source of light and view from the bottom. Rotate the comparator until the best color match is found (fig. 3).







Test Method

The Hydrazine CHEMets® and ULR CHEMets®1 test method employs the PDMAB chemistry.^{2,3} In an acidic solution, hydrazine reacts with PDMAB (p-dimethylaminobenzaldehyde) to form a yellow colored complex in direct proportion to the hydrazine concentration.

- CHEMets & ULR CHEMets are registered trademarks of CHEMetrics, Inc. U.S. Patent No. 3.634.038
- L. C. Thomas and G. J. Chamberlin, Colorimetric Chemical Analytical Methods. 8th ed., p. 195, Method I (1974)
- 3. ASTM D 1385 07, Hydrazine in Water

Visit www.chemetrics.com to view product demonstration videos.

Always follow the test procedure above to perform a test.



www.chemetrics.com 4295 Catlett Road, Midland, VA 22728 U.S.A. Phone: (800) 356-3072; Fax: (540) 788-4856 E-Mail: orders@chemetrics.com

Mar. 18, Rev. 6